TRIPLEMAX ITAPE

VETMED

ORAL DRENCH FOR SHEEP

Active Ingredients: 1g/L Ivermectin, 40g/L Levamisole HCL, 22.7g/L Oxfendazole, 18.8g/L Praziquantel with 0.5g/L Selenium & 2.2g/L Cobalt.

ADMINISTRATION

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

SHAKE WELL BEFORE USE.

VETMED Triplemax iTape is ready to use through standard drenching equipment. Rinse gun with water immediately after use. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing. Correct drenching technique should be used. Check accuracy of drench guns regularly.

DOSAGE GUIDE: Sheep by Oral Drench ImL/Skg body weight delivering 0.2mg/kg Ivermectin, 8mg/kg Levamisole HCL, 4.5mg/kg Oxlendazole, 3.76mg/kg Praziquantel, 0.1mg/kg Selenium and 0.44mg/kg Cobalt.

BODYWEIGHT	DOSE VOLUME	BODYWEIGHT	DOSE VOLUME
16 - 20kg	4 mL	46 - 50 kg	10 mL
21 - 25kg	5 mL	51 - 55 kg	11 mL
26 - 30 kg	6 mL	56 - 60 kg	12 mL
31 - 35 kg	7 mL	61 - 65 kg	13 mL
36 - 40 kg	8 mL	66 - 70 kg	14 mL
41 - 45 kg	9 mL	71 - 75 kg	15 mL
OVER 75KG ADD 1mL/5KG			

WITHHOLD PERIODS

MILK: Milk intended for sale for human consumption must be discarded during treatment and for 35 days following the last treatment.

MEAT: Sheep producing meat or offal for human consumption must not be sold for slaughter either during or within 14 days of the last treatment.

STORAGE

Store in a cool dry place at room temperature (i.e. 5° C to 25° C).

EFFICACY

VETMED Triplemax iTape Oral Drench contains Ivermectin; a member of the Macrocyclic Lactones, Levamisole; a member of the Imidazothiazole class of anthelmintics, Oxfendazole; a member of the Benzimidazoles and Praziquantel; a member of the Pyrozine family of anthelmintics. These actives are effective against internal parasites sensitive to their action family.

For the effective treatment and control of the following species of gastrointestinal parasites:

Gastrointestinal parasites:

For the treatment and control of the following mature and immature gastrointestinal parasites (*includes inhibited fourth stage larvae):

Ostertagia circumcincta*, Haemonchus contortus*, Trichostrongylus axei, Trichostrongylus colubriformis, Cooperia curticei*, Cooperia oncophora*, Cooperia punctata*, Nematodirus fillicollis, Nematodirus spathiger, Strongyloides papillosis, Oesophagostomum venulosum, Chabertia ovina, Trichuris ovis and Moniezia expansa.

Anthelmintic resistant strains:

VETMED Triplemax iTape Oral Drench may be effective in some populations of gastrointestinal worms with single or dual resistance to the Avermectin/Milbemycin, Benzimidazole or Levamisole/Morantel anthelmintic action families.

Lungworm:

Dictyocaulus filaria.

PACK SIZES

VETMED Triplemax iTape is available in 10L, 5L, and 1L packs.

VETMED Triplemax iTape is registered pursuant to the ACVM Act 1997 No. A11111.



